

Salmonella

Salmonella is extremely common in wild reptiles. It's also relatively common in wild birds, domestic poultry, and wild rodents. It's not a concern in laboratory rodents unless the animals have been deliberately infected as part of an experiment.

Salmonellosis can also be seen in horses and cattle; workers are at risk when treating or performing necropsies on infected large animals. Regular hand washing, and the use of appropriate personal protective equipment, such as coveralls, masks, and gloves, will minimize the chances of your becoming infected in these situations.

What is salmonellosis?

Salmonellosis is an infection with a bacteria called Salmonella. Most persons infected with Salmonella develop diarrhea, fever, and abdominal cramps 12 to 72 hours after infection. The illness usually lasts 4 to 7 days, and most persons recover without treatment. However, in some persons the diarrhea may be so severe that the patient needs to be hospitalized. In these patients, the Salmonella infection may spread from the intestines to the blood stream, and then to other body sites and can cause death unless the person is treated promptly with antibiotics. The elderly, infants, and those with impaired immune systems are more likely to have a severe illness.

What sort of germ is Salmonella?

The Salmonella germ is actually a group of bacteria that can cause diarrheal illness in humans. They are microscopic living creatures that pass from the feces of people or animals, to other people or other animals. There are many different kinds of Salmonella bacteria. Salmonella serotype Typhimurium and Salmonella serotype Enteritidis are the most common in the United States. Salmonella has been known to cause illness for over 100 years.

How can Salmonella infections be diagnosed?

Many different kinds of illnesses can cause diarrhea, fever, or abdominal cramps. Determining that Salmonella is the cause of the illness depends on laboratory tests that identify Salmonella in the stools of an infected person. These tests are sometimes not performed unless the laboratory is instructed specifically to look for the organism. Once Salmonella has been identified, further testing can determine its specific type, and which antibiotics could be used to treat it.

How can Salmonella infections be treated?

Salmonella infections usually resolve in 5-7 days and often do not require treatment unless the patient becomes severely dehydrated or the infection spreads from the intestines. Persons with severe diarrhea may require rehydration, often with intravenous fluids. Antibiotics are not usually necessary unless the infection spreads from the intestines, then it can be treated with ampicillin, gentamicin, trimethoprim/sulfamethoxazole, or ciprofloxacin. Unfortunately, some Salmonella bacteria have become resistant to antibiotics, largely as a result of the use of antibiotics to promote the growth of feed animals.

Are there long term consequences to a Salmonella infection?

Persons with diarrhea usually recover completely, although it may be several months before their bowel habits are entirely normal. A small number of persons who are infected with Salmonella, will go on to develop pains in their joints, irritation of the eyes, and painful urination. This is called Reiter's syndrome. It can last for months or years, and can lead to chronic arthritis which is difficult to treat. Antibiotic treatment does not make a difference in whether or not the person later develops arthritis.

How do people catch Salmonella?

Salmonella live in the intestinal tracts of humans and other animals, including birds. Salmonella are usually transmitted to humans by eating foods contaminated with animal feces. Contaminated foods usually look and smell normal. Contaminated foods are often of animal origin, such as beef, poultry, milk, or eggs, but all foods, including vegetables may become contaminated. Many raw foods of animal origin are frequently contaminated, but fortunately, thorough cooking kills Salmonella. Food may also become contaminated by the unwashed hands of an infected food handler, who forgot to wash his or her hands with soap after using the bathroom.

Salmonella may also be found in the feces of some pets, especially those with diarrhea, and people can become infected if they do not wash their hands after contact with these feces. Reptiles are particularly likely to harbor Salmonella and people should always wash their hands immediately after handling a reptile, even if the reptile is healthy

What can a person do to prevent this illness?

People should wash their hands after contact with animal feces. Since reptiles are particularly likely to have Salmonella, everyone should immediately wash their hands after handling reptiles.

How common is salmonellosis?

Every year, approximately 40,000 cases of salmonellosis are reported in the United States. Because many milder cases are not diagnosed or reported, the actual number of infections may be twenty or more times greater. Salmonellosis is more common in the summer than winter.

Children are the most likely to get salmonellosis. Young children, the elderly, and the immunocompromised are the most likely to have severe infections. It is estimated that approximately 1,000 persons die each year with acute salmonellosis.

Antibiotics as growth promotants in food animals. The US Department of Agriculture monitors the health of food animals, inspects egg pasteurization plants, and is responsible for the quality of slaughtered and processed meat. The US Environmental Protection Agency regulates and monitors the safety of our drinking water supplies.